

METHODS FOR IMPROVING CELL GROWTH AND ALCOHOL PRODUCTION DURING FERMENTATION

Related Information

- 5 This application claims priority to U.S. provisional application number 60/214,099 entitled "Stimulation of Growth and Ethanol Production in Engineered *Escherichia Coli* Resulting from the Addition of Acetaldehyde" filed June 26, 2000, and U.S. provisional application number 60/219,844 entitled "Methods for Improving Cell Growth and Alcohol Production During Fermentation" filed July 21, 2000, both of
- 10 which are incorporated herein in their entireties by this reference. The contents of all patents, patent applications, and references cited throughout this specification are hereby incorporated by reference in their entireties.

Government Sponsored Research

- 15 This work was supported, in part, by U.S. Dept. of Agriculture, National Research Initiative Grant No.: 98-35504-6177, and U.S. Dept. of Energy, Basic Energy Sciences, Grant No.: FG02-96ER20222.

Background of the Invention

- 20 Many environmental and societal benefits would result from the replacement of petroleum-based automotive fuels with renewable fuels obtained from plant materials (Lynd *et al.*, (1991) *Science* 251:1318-1323; Olson *et al.*, (1996) *Enzyme Microb. Technol.* 18:1-17; Wyman *et al.*, (1995) *Amer. Chem. Soc. Symp.* 618:272-290). Each year, the United States burns over 120 billion gallons of automotive fuel, roughly
- 25 equivalent to the total amount of imported petroleum. The development of ethanol as a renewable alternative fuel has the potential to eliminate United States dependence on imported oil, improve the environment, and provide new employment (Sheehan, (1994) ACS Symposium Series No. 566, ACS Press, pp 1-53).

- In theory, the solution to the problem of imported oil for automotive fuel appears
- 30 quite simple. Rather than using petroleum, a finite resource, the ethanol can be produced efficiently by the fermentation of plant material, a renewable resource. Indeed, Brazil has demonstrated the feasibility of producing ethanol and the use of ethanol as a primary automotive fuel for more than 20 years. Similarly, the United States produces over 1.2 billion gallons of fuel ethanol each year. Currently, fuel
- 35 ethanol is produced from corn starch or cane syrup utilizing either *Saccharomyces cerevisiae* or *Zymomonas mobilis* (*Z. mobilis*). However, both cane sugar and corn starch are relatively expensive starting materials, which have competing uses as food products.


Although some aspects of a biomass conversion process have been demonstrated, ethanol and other chemicals produced from biomass must be cost-competitive with existing petroleum-based products. These costs include nutrients and materials needed for bioconversion, production purification, waste treatment, power, and the manufacturing facility itself.

Therefore, methods that would reduce the cost associated with fermentation, including savings from a reduction in added nutrients and improvements in the rate of production and yield of product, *e.g.*, ethanol, would be beneficial.

10 Summary of the Invention


The present invention provides methods, which overcome the above stated problems of the high cost associated with the production of an alcohol, *e.g.*, ethanol, by fermentation. The invention provides a method for increasing the rate of alcohol production (*e.g.*, ethanol) and the growth of alcohologenic cells (*e.g.*, ethanologenic cells) by contacting or exposing such cells (*e.g.*, by culturing) with a nutrient compound (*e.g.*, a compound of formula I described below) which improves the productivity of the culture (*e.g.*, fermentation rate) and/or growth of the culture (*e.g.*, ability of the cells to grow to a higher cell density or having a reduced cell replication time).

More particularly, in a first aspect, the invention provides a method for increasing production of alcohol from a saccharide source by an alcohologenic cell by, contacting a saccharide source with an alcohologenic cell, and exposing the cell to at least one compound of formula I,



where R_1 is H, OH or COOR_2 ; R_2 is H or alkyl; R_3 is H, NH_2 , alkyl or alkenyl; R_4 is H, alkyl, alkenyl, or a side chain of a naturally occurring amino acid; and salts thereof; where the exposing results in the increased production of alcohol by the alcohologenic cell as compared to a control.

In a second aspect, the invention provides a method for increasing growth of a cell by, contacting a cell with a saccharide source, and exposing the cell to at least one compound of formula I,



where R_1 is H, OH or COOR₂; R_2 is H or alkyl; R_3 is H, NH₂, alkyl or alkenyl; R_4 is H, alkyl, alkenyl, or a side chain of a naturally occurring amino acid; and salts thereof; where the exposing results in the increased growth of the cell as compared to a control.

- 5 In one embodiment of the first two aspects, the compound of formula I is a lower aliphatic aldehyde, lower aliphatic α -keto carboxylic acids, lower aliphatic dicarboxylic acid, amino acid, or salt of any of the foregoing acids.

In one embodiment of the first aspect, the alcohol is ethanol and the alcoholicogenic cell is an ethanologenic cell. In a related embodiment, the increased
10 production of ethanol is indicated by an increase in volumetric productivity, preferably where the volumetric productivity is between about 0.3 g/L and about 0.5 g/L.

- In one embodiment of the above two aspects, the cell is selected from the family Enterobacteriaceae, more preferably, from the genus *Escherichia* or *Klebsiella*. In a related embodiment, the cell is *E. coli* KO4 (ATCC 55123), *E. coli* KO11 (ATCC
15 55124), *E. coli* KO12 (ATCC 55125), *K. oxytoca* M5A1, or *K. oxytoca* P2 (ATCC 55307), LY01 (ATCC _____). In another related embodiment, the cell is a recombinant cell.

- In another embodiment of the above aspects, the compound of formula I is acetaldehyde, pyruvate, succinate, isocitrate, glutamate, α -ketoglutarate, a yeast extract,
20 or casamino acids, and preferably, is acetaldehyde, pyruvate, or glutamate, α -ketoglutarate or a combination thereof.

In a related embodiment, the cell is exposed to glutamate and acetaldehyde, pyruvate and acetaldehyde, fumarate and malate, or α -ketoglutarate and succinate.

- In one embodiment, the cell is in an aqueous solution.

- 25 In even another embodiment, the saccharide source is cellooligosaccharide, lignocellulose, hemicellulose, cellulose, pectin, xylose, glucose, corn steep liquor (CSL), or any combination thereof.

In another embodiment, the cell is exposed to the compound of formula I for a period of time between about 1 and about 96 hours.

- 30 In another embodiment, the method is performed at a pH between about 6 and about 8, and preferably at a pH of about 6.5.

In another embodiment, the method is performed at a temperature between about 20° and about 40° C, and preferably at a temperature of about 35° C.

- 35 In another embodiment, the compound is present at a concentration between about 0.1 and about 4.0 g/L.

In another embodiment, the method of the above aspects further includes exposing the cell to the compound more than once. In a related embodiment, the

exposing of the cell to the compound is performed at time intervals between about 1 hour and about 24 hours.

In another embodiment, the method of the above aspects further includes exposing the cell to two or more different compounds of formula I.


- 5 In another embodiment, the method of the above aspects further includes agitating the cell, the saccharide source, and the compound between about 50 rpm and about 200 rpm.

- In one embodiment of the second aspect, the increased growth is indicated by increased cell density or decreased cell replication time. In a related embodiment, the increased cell density is indicated by an optical density of between about 2 and about 3 at 550 nm after 24 hours.

- In another embodiment, the method of the above aspects is performed in a fermentor vessel, where, preferably, the cell and the saccharide source are provided in an aqueous solution. In a related embodiment, the aqueous solution includes a fermentation medium, preferably Luria broth or CSL broth.

In yet another embodiment, the method of the above aspects is suitable for simultaneous saccharification and fermentation.

- In a third aspect, the present invention provides a growth medium suitable for use in an improved fermentation process including a saccharide source, a basal nutrient medium, and at least one compound of formula I,



where R₁ is H, OH or COOR₂; R₂ is H or alkyl; R₃ is H, NH₂, alkyl or alkenyl; R₄ is H, alkyl, alkenyl; or a side chain of a naturally occurring amino acid; and salts thereof.

- 30 In one embodiment, the saccharide source is cellooligosaccharide, lignocellulose, hemicellulose, cellulose, pectin, xylose, glucose, or any combination thereof.


In another embodiment, the basal nutrient medium is Luria broth or CSL broth.

In even another embodiment, the medium is suitable for use in simultaneous saccharification and fermentation.

- 35 In still another embodiment, the compound of formula I is acetaldehyde, pyruvate, succinate, citrate, isocitrate, glutamate, α-ketoglutarate, malate, fumarate, a yeast extract, or a casamino acid. In a related embodiment, the compound of formula I is preferably acetaldehyde, pyruvate, succinate, citrate, isocitrate, glutamate, α-ketoglutarate, or malate.

In even another embodiment, the growth medium is packaged with instructions for use.

In a fourth aspect, the invention provides a fermentation reaction mixture suitable for producing ethanol containing, a growth medium having a saccharide source, an ethanologenic cell, and an exogenous source of at least one compound of formula I,



where R_1 is H, OH or COOR_2 ; R_2 is H or alkyl; R_3 is H, NH_2 , alkyl or alkenyl; R_4 is H, alkyl, alkenyl; or a side chain of a naturally occurring amino acid; and salts thereof.

In one embodiment, the fermentation reaction mixture includes a saccharide source selected from the group consisting of cellooligosaccharide, lignocellulose, hemicellulose, cellulose, pectin, xylose, glucose, or any combination thereof.

In another embodiment, the ethanologenic cell of the fermentation reaction mixture is from the family Enterobacteriaceae.

In another embodiment, the fermentation reaction mixture is suitable for use in simultaneous saccharification and fermentation.

In still another embodiment, the compound of formula I is acetaldehyde, pyruvate, succinate, isocitrate, glutamate, α -ketoglutarate, fumarate, a yeast extract, or a casamino acid, and preferably, acetaldehyde, pyruvate, succinate, isocitrate, glutamate, or α -ketoglutarate.

Advantages of the above compositions and methods include the ability to reduce the overall cost of biomass conversion to a useable fuel. For example, increases in alcohol yield or alcohol titer provide the benefits of reducing the amount of biomass which must be treated accompanied by corresponding reductions in the costs of feedstocks, chemicals, equipment, and energy throughout the process. Improvements in yield and titer also reduces the amount of waste generated and the costs associated with waste disposal.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1 shows ethanol production and cell growth by ethanologenic bacteria when cultured in broth containing 1% corn steep liquor (CSL), xylose, salts, and

different amounts of an additional nutrient compound (*i.e.*, acetaldehyde) as compared to a control. Panel A shows ethanol production in g/L over time (96 hours), and panel B shows changes in cell growth (measured as cell mass at OD_{550nm}) over time (96 hours). Ethanol production and cell mass are determined at several time points from the start of fermentation to the end of the 96 hour time period. The additional nutrient compound acetaldehyde was added to the fermentation medium at five different concentrations and at five different time points as indicated.

Figure 2 shows ethanol production and cell growth by ethanologenic bacteria when cultured in broth containing 1% corn steep liquor (CSL), glucose, salts, and different amounts of an additional nutrient compound (*i.e.*, acetaldehyde) as compared to a control. Panel A shows ethanol production in g/L over time (96 hours), and panel B shows changes in cell growth (measured as cell mass at OD_{550nm}) over time (96 hours). Ethanol production and cell mass are determined at several time points from the start of fermentation to the end of the 72 hour time period. The additional nutrient compound acetaldehyde was added to the fermentation medium at five different concentrations and at five different time points as indicated.

Figure 3 shows ethanol production and cell growth by ethanologenic bacteria when cultured in Luria broth containing xylose and different amounts of an additional nutrient compound (*i.e.*, acetaldehyde) as compared to a control. Panel A shows ethanol production in g/L over time (72 hours), and panel B shows changes in cell growth (measured as cell mass at OD_{550nm}) over time (72 hours). Ethanol production and cell mass are determined at several time points from the start of fermentation to the end of the 72 hour time period. The additional nutrient compound acetaldehyde was added to the fermentation medium at five different concentrations and at five different time points as indicated.

Figure 4 is a schematic representation of the sugar to ethanol pathway indicating acetaldehyde as an intermediate metabolite in the pyruvate to ethanol pathway.

Figure 5 is a schematic representation of the overall fermentation pathway for hexoses and pentoses.

Detailed Description of the Invention

Before further description of the invention, certain terms employed in the specification, examples and appended claims are, for convenience, collected here.

I. Definitions

As used herein, the term “medium” or “media”, refers to an aqueous or solid source of nutrients capable of supporting the growth of a cell, preferably, for example, an alcohologenic cell capable of fermenting a carbon source, such as a sugar, into an

alcohol. Examples of media include, *e.g.*, Luria broth (LB), NZCYM medium, NZYM medium, NZM medium, SOB medium, SOC medium, ZXYT medium, M9 minimal medium, Terrific broth (TB) (see also, Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, CSHL Press (1989); Ausubel *et al.*, Current Protocols in Molecular

5 Biology, Wiley Interscience).

The term "Luria broth" or "LB" includes media typically comprising a yeast extract (*e.g.*, crude, self digested solubles from yeast bodies containing, *e.g.*, amino acids, peptides, vitamins, lipids, nucleosides, salts, *etc.*), casamino acids (*i.e.*, an enzymatic digestion of casein protein comprising amino acids and peptides), and salts

10 (*e.g.*, sodium chloride).


The term "CSL medium" includes a medium typically comprising corn steep liquor, a fermentable sugar, and a mixture of salts essential for growth.

The term "cell", refers to the smallest structure capable of independently carrying out life sustaining processes, including metabolic processes, *e.g.*, growth, and reproduction. The term "cell," as used herein, includes a bacterial, yeast, fungal, plant,

15 or animal cell.

The term "nutrient compound", includes any molecule or compound, added to the medium of a cell during the culture of the cell for the purpose of improving product yield or cell growth as compared to a control. The term includes any compound of the

20 formula I:



25

wherein;

R₁ is H, OH or COOR₂;

R₂ is H or alkyl;

30 R₃ is H, NH₂, alkyl or alkenyl;

R₄ is H, alkyl, alkenyl, or a side chain of a naturally occurring amino acid, and salts thereof. Preferred nutrient compounds of formula I include but are not limited to, lower aliphatic aldehydes, lower aliphatic α -keto carboxylic acids, lower aliphatic dicarboxylic acids, amino acids, and salts of any of these acids. Preferably,

35 carboxylic acid compounds of formula I are used as salts, *e.g.*, mono- or bi- potassium and/or sodium salts, hydrated or unhydrated. Particularly preferred compounds of formula I are those listed in Tables 1-7, including but not limited to, acetaldehyde, pyruvate, glutamate, aspartate, isocitrate, oxaloacetate, alanine, succinate, fumarate,

malate, α -ketoglutarate, yeast extract, and amino acids, *e.g.*, casamino acids, separately or in any combination.

The term "alkyl" is art-recognized and includes the radical of saturated aliphatic groups, including straight-chain alkyl groups, branched-chain alkyl groups, cycloalkyl (alicyclic) groups, alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In preferred embodiments, a straight chain or branched chain alkyl has 30 or fewer carbon atoms in its backbone (*e.g.*, C₁-C₃₀ for straight chain, C₃-C₃₀ for branched chain), more preferably 20 or fewer, and still more preferably four or fewer. Likewise, preferred cycloalkyls have from four to ten carbon atoms in their ring structure, and more preferably have 5, 6, or 7 carbons in the ring structure.

Unless the number of carbons is otherwise specified, the term "lower" as in "lower alkyl" and/or "lower aliphatic" is intended to denote a saturated or unsaturated aliphatic hydrocarbon (*e.g.*, alkyl or alkenyl as defined herein) having from one to ten carbons, more preferably from one to six, and most preferably from one to four carbon atoms in its backbone structure, which may be straight or branched-chain. Examples of lower alkyl groups include methyl, ethyl, n-propyl, i-propyl, tert-butyl, hexyl, heptyl, octyl, and so forth. Likewise, "lower alkenyl" and "lower alkynyl" have similar chain lengths. Preferred alkyl groups include lower alkyls. Examples of alkylene groups are methylene, ethylene, propylene, and so forth.

Moreover, the term alkyl as herein is intended to include both "unsubstituted alkyls" and "substituted alkyls", the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, halogen, hydroxyl, carbonyl (including aldehydes, ketones, carboxylates, and esters), alkoxyl, ether, phosphoryl, cyano, amino, acylamino, amido, amidino, imino, sulfhydryl, alkylthio, arylthio, thiolcarbonyl (including thioformates, thiolcarboxylic acids, and thioesters), sulfonyl, nitro, heterocyclyl, aralkyl, or an aromatic or heteroaromatic moiety. It will be understood by those skilled in the art that the moieties substituted on the hydrocarbon chain can themselves be substituted, if appropriate. For instance, the substituents of a substituted alkyl may include substituted and unsubstituted forms of amino, acylaminos, iminos, amidos, phosphoryls (including phosphonates and phosphinates), sulfonyls (including sulfates, sulfonates, sulfamoyls, and sulfonamidos), and silyl groups, as well as ethers, alkylthios, arylthios, carbonyls (including ketones, aldehydes, carboxylates, and esters), -CF₃, -CN, and the like. Exemplary substituted alkyls are described below. Cycloalkyls can be further substituted with alkyls, alkenyls, alkoxys, alkylthios, arylthios, aminoalkyls, carbonyl-substituted alkyls, -CF₃, cyano (-CN), and the like.

The term "aralkyl", as used herein, refers to an alkyl group substituted with an aryl group (*e.g.*, an aromatic or heteroaromatic group).

The terms "alkenyl" and "alkynyl" are art-recognized and include unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond, respectively.

The term "alkoxyl" is art-recognized and includes any group represented by the formula -O-alkyl. Representative alkoxyl groups include methoxy, ethoxy, propoxy, tert-butoxy, and the like. Unless otherwise specified, an "alkoxy" group can be replaced with a group represented by -O-alkenyl, -O-alkynyl, -O-aryl (*i.e.*, an aryloxy group), or -O-heterocyclyl. An "ether" is two substituted or unsubstituted hydrocarbons covalently linked by oxygen. Accordingly, the substituent of, *e.g.*, an alkyl that renders that alkyl an ether is, or resembles, an alkoxyl, such as can be represented by one of -O-alkyl, -O-alkenyl, -O-alkynyl, -O-aryl, or -O-heterocyclyl. The term "lower alkoxy" includes a lower alkyl group attached to the remainder of the molecule by oxygen.


Examples of alkoxy groups include methoxy, ethoxy, isopropoxy, tert-butoxy and so forth. The term "phenyl alkoxy" refer to an alkoxy group, which is substituted by a phenyl ring. Examples of phenyl alkoxy groups are benzyloxy, 2-phenylethoxy, 4-phenylbutoxy, and so forth. The term "alkanoyloxy group" refers to the residue of an alkylcarboxylic acid formed by removal of the hydrogen from the hydroxyl portion of the carboxyl group. Examples of alkanoyloxy groups include formyloxy, acetoxy, butyryloxy, hexanoyloxy, and so forth. The term "substituted" as applied to "phenyl" refers to phenyl which is substituted with one or more of the following groups: alkyl, halogen (*i.e.*, fluorine, chlorine, bromine or iodine), nitro, cyano, trifluoromethyl, and so forth. The "alkanol" or a "hydroxyalkyl" refer to a compound derived by protonation of the oxygen atom of an alkoxy group. Examples of alkanols include methanol, ethanol, 2-propanol, 2-methyl-2-propanol, and the like.

The term "halogen" designates -F, -Cl, -Br or -I; the term "sulfhydryl" or "thiol" means -SH; the term "hydroxyl" means -OH.

The term "aryl" is art-recognized and includes 5- and 6-membered single-ring aromatic groups that may include from zero to four heteroatoms, for example, benzene, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, pyrazole, pyridine, pyrazine, pyridazine and pyrimidine, and the like. Aryl groups also include polycyclic fused aromatic groups such as naphthyl, quinolyl, indolyl, and the like. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles", "heteroaryls", or "heteroaromatics". The aromatic ring can be substituted at one or more ring positions with such substituents as described above, as for example, halogen, alkyl, aralkyl, alkenyl, alkynyl, cycloalkyl, hydroxyl, amino, acylamino, azido, nitro, sulfhydryl, imino, amido, amidino, phosphonate, phosphinate, carbonyl, carboxyl, silyl, ether, alkylthio, arylthio, sulfonyl, sulfonamido, sulfamoyl, ketone, aldehyde, ester, a heterocyclyl, an aromatic or heteroaromatic moiety, -CF₃, -CN, or the like: Aryl

groups can also be fused or bridged with alicyclic or heterocyclic rings, which are not aromatic so as to form a polycycle (e.g., tetralin).

The term "amino acid" includes its art recognized meaning and broadly encompasses compounds of formula II:



Preferred amino acids include the naturally occurring amino acids, as well as synthetic derivatives, and amino acids derived from proteins, e.g., proteins such as casein, i.e., casamino acids, or enzymatic or chemical digests of, e.g., yeast, an animal product, e.g., a meat digest, or a plant product, e.g., soy protein, cottonseed protein, or a corn steep liquor (see, e.g., Traders' Guide to Fermentation Media, Traders Protein, Memphis, TN (1988), Biotechnology: A Textbook of Industrial Microbiology, Sinauer Associates, Sunderland, MA (1989), and Product Data Sheet for Corn Steep Liquor, Grain Processing Corp., IO).

The term "naturally occurring amino acid" includes any of the 20 amino acid residues which commonly comprise most polypeptides in living systems, rarer amino acids found in fibrous proteins (e.g., 4-hydroxyproline, 5-hydroxylysine, ε-N-methyllysine, 3-methylhistidine, desmosine, isodesmosine), and naturally occurring amino acids not found in proteins (e.g., β-alanine, γ-aminobutyric acid, homocysteine, homoserine, citrulline, ornithine, canavanine, djenkolic acid, and β-cyanoalanine).

The term "side chain of a naturally occurring amino acid" is intended to include the side chain of any of the naturally occurring amino acids, as represented by R in formula II. One skilled in the art will understand that the structure of formula II is intended to encompass amino acids such as proline where the side chain is a cyclic or heterocyclic structure (e.g., in proline R group and the amino group form a five-membered heterocyclic ring. Similarly, the compound of formula I above is intended to encompass amino acids such as proline wherein in formula I, e.g., R3 and R4 form a heterocyclic ring.

The term "increasing production of alcohol" refers to any increase in the yield of alcohol, e.g., ethanol, the volumetric productivity of a fermentation reaction, or the rate of production of alcohol, e.g., ethanol, from a fermentation reaction over a certain period of time or at the completion of the fermentation reaction, as compared to a control.

The term "volumetric productivity" includes the increased productivity of a cell culture where the productivity of the cells is typically measured as an increase in the

amount of a cell derived product in a given cell culture volume, preferably, *e.g.*, an increase in the amount of alcohol produced in grams per liter of culture (*i.e.*, g/L).

The term "increasing growth of a cell" includes increased cell density or cell mass, and/or decreased cell replication time as compared to a control. Cell mass and cell density may be determined by the optical density (OD) of the cells in suspension at any given time point.

The term "exposing" includes contacting the cell with a nutrient compound, *e.g.*, acetylaldehyde from any source. The cell may or may not be in aqueous solution.

The term "fermentation reaction" refers to any mixture of medium and cells capable of fermenting a saccharide source.

The term "fermentor vessel" refers to any container capable of supporting a fermentation reaction. A fermentor vessel may be capable of containing a volume of between 0.10 to 100L, or more (*e.g.*, 1,000,000 L). A fermentor vessel may also have a means of controlling temperature and pH and may provide a source of agitation (*e.g.*, via an impeller and/or sparging) for the contents of the vessel. In addition, a fermentor vessel may also provide a source of gas flow (oxygen, nitrogen, and/or carbon dioxide). Typically a fermentor vessel allows for all or some of the foregoing culture characteristics or parameters to be advantageously monitored and/or controlled.

The term "exogenous source" is intended to include any source of a nutrient compound that is added to the fermentation reaction. Examples of exogenous sources of nutrient compounds are described in the Examples.

The term "basal nutrient medium" is any medium, which contains all of the elements essential for maintaining the fundamental vital activities of an organism. For example, a basal nutrient medium includes Luria broth (LB).

The term "recombinant cell" is intended to include a genetically modified cell. The cell can be a microorganism or a higher eukaryotic cell. The term is intended to include progeny of the cell originally modified. In preferred embodiments, the cell is a alcohologenic bacterial cell, *e.g.*, a Gram-negative bacterial cell, and this term is intended to include all facultatively anaerobic Gram-negative cells of the family Enterobacteriaceae such as *Escherichia*, *Shigella*, *Citrobacter*, *Salmonella*, *Klebsiella*, *Enterobacter*, *Erwinia*, *Kluyvera*, *Serratia*, *Cedecea*, *Morganella*, *Hafnia*, *Edwardsiella*, *Providencia*, *Proteus*, and *Yersinia*. Particularly preferred recombinant hosts are *Escherichia coli* or *Klebsiella oxytoca* cells having alcohologenic activities. More preferred host cells have polysaccharase and alcohologenic activities and can ferment a complex sugar. Examples of such cells are provided in U.S. Patent Nos. 5,821,093; 5,482,846; 5,424,202; 5,028,539; 5,000,000; 5,487,989, 5,554,520; 5,162,516; and USSN 60/136,376.

The term "polysaccharase" includes a polypeptide capable of catalyzing the degradation or depolymerization of any linked sugar moiety, *e.g.*, disaccharides, trisaccharides, oligosaccharides, including, complex carbohydrates, *i.e.*, complex sugars, *e.g.*, lignocellulose, which comprises cellulose, hemicellulose, and pectin. The terms are intended to include cellulases such as glucanases, including both endoglucanases and exoglucanases, and β -glucosidase.

The term "complex sugar" includes any carbohydrate source comprising more than one sugar molecule. These carbohydrates may be derived from any unprocessed plant material or any processed plant material. Examples are wood, paper, pulp, plant derived fiber, or synthetic fiber comprising more than one linked carbohydrate moiety, *i.e.*, one sugar residue. One particular complex sugar is lignocellulose, which represents approximately 90% of the dry weight of most plant material and contains carbohydrates, *e.g.*, cellulose, hemicellulose, pectin, and aromatic polymers, *e.g.*, lignin. Cellulose makes up 30%-50% of the dry weight of lignocellulose and is a homopolymer of cellobiose (a dimer of glucose).

The term "saccharide source" includes any sugar including, for example, monosaccharides, disaccharides, oligosaccharides, complex sugars, or any combination thereof. Exemplary saccharide sources include, *e.g.*, glucose and xylose. Any one or a combination of the above carbohydrates are potential sources of sugars for depolymerization (if needed) and subsequent bioconversion to an alcohol, *e.g.*, ethanol, by fermentation according to the present invention.

The term "simultaneous saccharification and fermentation" or "SSF" is intended to include the use of one or more cells, *e.g.*, recombinant cells, for the contemporaneous degradation or depolymerization of a complex sugar and bioconversion of that sugar into an alcohol, *e.g.*, ethanol, by fermentation.

The term "ethanologenic" is intended to include the ability of a microorganism to produce ethanol from a carbohydrate as a primary fermentation product. The term is intended to include naturally occurring ethanologenic organisms, organisms with naturally occurring or induced mutations, and organisms which have been genetically modified.

The term "Gram-negative bacteria" is intended to include the art recognized definition of this term. Typically, Gram-negative bacteria include, for example, the family Enterobacteriaceae which comprises, among others, the species *Escherichia* and *Klebsiella*.

The term "alcohologenic" includes the ability of a cell, preferably of a microorganism, to produce an alcohol, *e.g.*, a carbon-based molecule with a hydroxyl moiety, *e.g.*, ethanol, from a carbohydrate as a primary fermentation product. The term is intended to include naturally occurring alcohologenic organisms, organisms with

naturally occurring or induced mutations, and organisms which have been genetically modified.

The term "alcohol" refers to any carbon based molecule having a hydroxyl group such as, *e.g.*, ethanol, but also including, *e.g.*, methanol, propanol, butanol, *etc.*

The term "control" includes its art recognized meaning and, *e.g.*, typically refers to a sample or culture exposed to the same conditions as the test culture but for one parameter such as, *e.g.*, an additional nutrient compound in the medium; *i.e.*, the control sample would not contain the additional nutrient compound, preferably, *e.g.*, a compound represented by formula I, *supra*.

The term "carbon-based energy source", "sugar", or "saccharide source" are used interchangeably and include any sugar that can be metabolized by a cell.

II. Increased Ethanol Production and Cell Growth

The present invention relates, in part, to a method for increasing the rate of alcohol, *i.e.*, ethanol, production and final ethanol titer from a saccharide source by the addition of one or more compounds to fermenting cultures of alcohologenic cells (*e.g.*, ethanologenic cells), as compared to a control with no additional compound. A compound is any of the compounds listed in Tables 1-7, in the Examples. For example, in a preferred embodiment, acetaldehyde and pyruvate are compounds of the invention.

Other examples of compounds include, but are not limited to glutamate, aspartate, isocitrate, oxaloacetate, alanine, succinate, fumarate, malate, α -ketoglutarate, yeast extract (an amino source as well as vitamins, minerals, lipids, *etc.*), and casamino acids (amino acids derived from casein). Compounds of the methods of the invention may be added separately or in any combination.

Fermentation products such as ethanol are essentially waste products of sugar metabolism, essential for electron balance and the regeneration of NAD⁺. Acetaldehyde is a product of ethanol-producing microorganisms and an intermediate metabolite in the pyruvate to ethanol pathway (see Fig. 4, and Fig. 5 for a more general schematic). It is produced by the non-oxidative decarboxylation of pyruvate by pyruvate decarboxylase, and subsequently reduced to ethanol during the oxidation of NADH by alcohol dehydrogenase.

Moreover, the present invention relates, in part, to a method of increasing the rate of growth and the final cell concentration achieved in fermenting cultures of cells by the addition of one or more compound listed in Tables 1-7 to the cell culture.

Accordingly, an increase in cell growth leads to a higher rate of ethanol production per unit volume during fermentation. Increased growth of the cells may be determined by increased cell density and/or decreased cell replication time. The increase in cell density over time can be used to measure the growth of the cells in culture. Cell mass can be

determined by the optical density (OD) of the cells at any given time point. The maximum cell density is the time at which the cell culture has reached the maximum OD. In one embodiment, increased cell density is determined when the cell density is between an optical density of 2 and 3 at 550 nm.

- 5 Gas chromatography is advantageously used to measure the increase of ethanol production after addition of a compound as compared to a control. In one embodiment, the production of ethanol is an increase in volumetric productivity. In a preferred embodiment, volumetric productivity is between 0.3 and 0.5 g/L.

- In another embodiment of the invention, the method of the invention is
10 performed in a fermentor vessel, allowing for larger volumes of ethanol production from a reduced number of fermentation vessels, and a further reduction of cost in the bioconversion process. A fermentor vessel, as used herein, is any vessel capable of supporting a fermentation reaction.

- In one embodiment, the cell used in the methods of the present invention is
15 selected from the family Enterobacteriaceae. For example, the cell may be an *Escherichia* or a *Klebsiella* cell. Exemplary *E. coli* strains that are ethanologenic include, for example, KO4 (ATCC 55123), KO11 (ATCC 55124), and KO12 (ATCC 55125) strains, as well as the LY01 (ATCC _____) strain, an ethanol-tolerant mutant of the *E. coli* strain KO11. Ideally, these strains may be derived from the *E. coli*
20 strain ATCC 11303, which is hardy to environmental stresses and can be engineered to be ethanologenic and secrete a polysaccharase/s. In addition, recent PCR investigations have confirmed that the ATCC 11303 strain lacks all genes known to be associated with the pathogenicity of *E. coli* (Kuhnert *et al.*, (1997) *Appl. Environ. Microbiol.* 63:703-709).

- A preferred ethanologenic bacterium is the *E. coli* KO11 strain which is capable
25 of fermenting hemicellulose hydrolysates from many different lignocellulosic materials and other substrates (Asghari *et al.*, (1996) *J. Ind. Microbiol.* 16:42-47; Barbosa *et al.*, (1992) *Current Microbiol.* 28:279-282; Beall *et al.*, (1991) *Biotechnol. Bioeng.* 38:296-303; Beall *et al.*, (1992) *Biotechnol. Lett.* 14:857-862; Hahn-Hagerdal *et al.*, (1994)
30 *Appl. Microbiol. Biotechnol.* 41:62-72; Moniruzzaman *et al.*, (1996) *Biotechnol. Lett.* 18:955-990; Moniruzzaman *et al.*, (1998) *Biotechnol. Lett.* 20:943-947; Grohmann *et al.*, (1994) *Biotechnol. Lett.* 16:281-286; Guimaraes *et al.*, (1992) *Biotechnol. Bioeng.* 40:41-45; Guimaraes *et al.*, (1992) *Biotechnol. Lett.* 14:415-420; Moniruzzaman *et al.*, (1997) *J. Bacteriol.* 179:1880-1886). This strain is able to rapidly ferment a
35 hemicellulose hydrolysate from rice hulls (which contained 58.5 g/L of pentose sugars and 37 g/L of hexose sugars) into ethanol (Moniruzzaman *et al.*, (1998) *Biotechnol. Lett.* 20:943-947). It was noted that this strain was capable of fermenting a hemicellulose

hydrolysate to completion within 48 to 72 hours, and under ideal conditions, within 24 hours.

Another preferred cell used in the methods of the present invention is the bacterium *Klebsiella*. In particular, *Klebsiella oxytoca* is preferred because, like *E. coli*, this enteric bacterium has the native ability to metabolize monomeric sugars, which are the constituents of more complex sugars. Moreover, *K. oxytoca* has the added advantage of being able to transport and metabolize cellobiose and cellotriose, the soluble intermediates from the enzymatic hydrolysis of cellulose (Lai *et al.*, (1996) *Appl. Environ. Microbiol.* 63:355-363; Moniruzzaman *et al.*, (1997) *Appl. Environ. Microbiol.* 63:4633-4637; Wood *et al.*, (1992) *Appl. Environ. Microbiol.* 58:2103-2110).

In one embodiment, the cell used in the methods of the present invention is a recombinant cell. Accordingly, the methods of the invention provide for use of genetically engineered ethanologenic derivatives of *K. oxytoca*, e.g., strain M5A1 having the *Z. mobilis* *pdg* and *adhB* genes encoded within the PET operon (as described in U.S.P.N. 5,821,093; Wood *et al.*, (1992) *Appl. Environ. Microbiol.* 58:2103-2110). The resulting organism, *K. oxytoca* P2 (ATCC 55307), produces ethanol efficiently from monomer sugars and from a variety of saccharides including raffinose, stachyose, sucrose, cellobiose, cellotriose, xylobiose, xylotriose, maltose, *etc.* (Burchhardt *et al.*, (1992) *Appl. Environ. Microbiol.* 58:1128-1133; Moniruzzaman *et al.*, (1997) *Appl. Environ. Microbiol.* 63:4633-4637; Moniruzzaman *et al.*, (1997) *J. Bacteriol.* 179:1880-1886; Wood *et al.*, (1992) *Appl. Environ. Microbiol.* 58:2103-2110).

In one embodiment, the methods of the present invention are suitable for simultaneous saccharification and fermentation (SSF). SSF is a process in which one or more recombinant hosts are used for the contemporaneous degradation or depolymerization of a complex sugar and bioconversion of that sugar residue into ethanol by fermentation. For example, the strain *K. oxytoca* P2 is suitable for use in the bioconversion of a complex saccharide in an SSF process as it contains polysaccharase genes in addition to ethanologenic activity (Doran *et al.*, (1993) *Biotechnol. Progress.* 9:533-538; Doran *et al.*, (1994) *Biotechnol. Bioeng.* 44:240-247; Wood *et al.*, (1992) *Appl. Environ. Microbiol.* 58:2103-2110). In particular, the use of this ethanologenic P2 strain eliminates the need to add supplemental cellobiase, one of the least stable components of commercial fungal cellulases (Grohmann, (1994) *Biotechnol. Lett.* 16:281-286). The addition of a nutrient compound to a SSF reaction, according to the methods of the invention, increases the production of ethanol during the simultaneous saccharification and fermentation process.

In another embodiment, the recombinant cell contains a polynucleotide segment that encodes a polysaccharase that is a glucanase, endoglucanase, exoglucanase, cellobiohydrolase, α -glucosidase, endo-1,4- α -xylanase, β -xylosidase, β -glucuronidase,

α -L-arabinofuranosidase, acylesterase, acetylxylosterase, α -amylase, β -amylase, glucoamylase, pullulanase, β -glucanase, hemicellulase, arabinosidase, mannanase, pectin hydrolase, pectate lyase, or a combination of these polysaccharases. In a related embodiment, the polysaccharase is a glucanase, preferably an expression product of

5 *celZ* or *celY* gene, and more preferably, derived from *Erwinia chrysanthemi*.

In one embodiment, the ethanologenic cells are exposed to a compound in an aqueous solution. For example, the fermentation media may be aqueous Luria broth (LB), variations thereof, other suitable medias, e.g., 1% CSL (see also those media described in the Examples) or media described in, e.g., Sambrook *et al.* or Ausubel *et*

10 *al.*, *supra*.

In another embodiment, the saccharide source from which ethanol is produced by the methods of the present invention is selected from the group consisting of celooligosaccharide, lignocellulose, hemicellulose, cellulose, pectin, xylose, glucose, corn steep liquor, and any combination thereof.

15 In one aspect of the invention, the method of exposing fermenting ethanologenic cells to a nutrient compound is performed over a period of time. In a preferred embodiment, the period of time is between about 1 hour and about 96 hours. The exposure of fermenting ethanologenic bacteria to a compound may be an exposure at one time point only, at more than one time point, or continuously. In one embodiment,

20 the exposure of a compound may be at several time points over a specific time period. For example, the compound may be added to the fermentation media, 1) at the time of inoculation of the fermentation media by an ethanologenic cell, 2) at the time of inoculation followed by a second addition after either 8 or 12 hours of fermentation, or 3) at the time of inoculation followed by subsequent additions after 12 hours and 24

25 hours. The addition of the compound may be at any time point during the fermentation of the saccharide source.

In one embodiment, the nutrient compound is, e.g., acetaldehyde or pyruvic acid, and is added to a final concentration between about 0.1 and about 4.0 g/L. The nutrient compound is preferably of the formula $R_3 - C(=O) - R_1$ where R_1 is H, OH or $COOR_2$,

30 R_2 is H or C_1-C_5 alkyl, R_3 is C_1-C_5 alkyl, or C_1-C_5 alkenyl and therefore includes, acetaldehyde, pyruvate, succinate, citrate, isocitrate, glutamate, α -ketoglutarate, malate, a casamino acid, a yeast extract, or any combination thereof (including free powder forms and salts thereof). Concentrations intermediate to the ranges cited above are also intended to be within the scope of the present invention (*i.e.*, 0.15 g/L, 0.2 g/L, 0.25 g/L,

35 0.3 g/L, 0.35 g/L, 0.4 g/L, 0.45 g/L, 0.5 g/L, 0.55 g/L, 0.6 g/L, 0.65 g/L, 0.7 g/L, 0.8 g/L, 0.9 g/L, 1.0 g/L, 1.5 g/L, 2.0 g/L, 2.5 g/L, 3.0 g/L, 3.5 g/L, and 4.0 g/L). It will be appreciated that no more than routine experimentation is needed for determining or optimizing, using the methods disclosed herein, a concentration or concentration range

for a given nutrient compound or combination of compounds. In a preferred embodiment, acetaldehyde may be added to a concentration of 0.1 g/L, 0.25 g/L, .5 g/L, or 0.75 g/L, or any combination thereof, during the fermentation process in order to achieve optimum production of ethanol and increased cell growth. To determine

- 5 increased production of ethanol and increased growth of the cell by the addition of a compound, distilled water may be added instead a compound as a control.

In another embodiment, the method of increasing production of ethanol and growth of the ethanologenic cell by the addition of a nutrient compound to fermenting ethanologenic bacteria can be performed at a pH between 6 and 8. pH values
10 intermediate to the ranges cited above are also intended to be within the scope of the present invention (e.g., 6.5, 7, and 7.5). In a preferred embodiment, the method of the instant invention is performed at a pH of about 6.5. The addition of a base, e.g., 2N KOH, to the fermentation medium can be used to maintain a specific pH during fermentation, as described in the examples.

- 15 Exposing a culture of fermenting cells to acetaldehyde can be performed at a temperature between about 20°C and about 40° C. Temperatures intermediate to the ranges cited above are also intended to be within the scope of the present invention (e.g., 21°C, 22°C, 23°C, 24°C, 25°C, 26°C, 27°C, 28°C, 29°C, 30°C, 31°C, 32°C, 33°C, 34°C, 35°C, 36°C, 37°C, 38°C, and 39°C). In a preferred embodiment, the method is
20 performed at a temperature of about 35°C. Fermentation may be performed with agitation between about 50 and about 200 rpm of the fermentation medium after inoculation with the cells and exposure to a compound. Rates of agitation intermediate to the ranges cited above are also intended to be within the scope of the present invention (i.e., 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, and 190).

- 25 The present invention is also based, in part, on a fermentation reaction suitable for producing alcohol, e.g., ethanol. In one embodiment, the growth medium has a saccharide source, an ethanologenic cell, and an exogenous source of a compound, where the fermentation reaction is incubated under conditions sufficient for producing ethanol. For example, the saccharide source of the fermentation reaction can be selected
30 from the group consisting of cellooligosaccharide, lignocellulose, hemicellulose, cellulose, pectin, xylose, glucose, and any combination thereof. An exogenous source of a compound may be any source obtained from outside of the fermentation reaction itself. The claimed fermentation reaction is also suitable for use in simultaneous saccharification and fermentation, where the degradation or depolymerization of a
35 complex sugar and bioconversion of that sugar residue into ethanol by fermentation takes place contemporaneously in a single fermentation reaction of the present invention.

Furthermore, another aspect of this invention includes a growth medium suitable for use in an improved fermentation reaction. The growth medium contains a saccharide source as described above, a basal nutrient medium such as, for example, Luria broth or 1% CSL, minerals, and a nutrient compound. The growth medium of the invention is suitable for use in simultaneous saccharification and fermentation.

III. Potential Substrates for Bioconversion into Ethanol

One advantage of the invention is the ability to use a saccharide source that has been, heretofore, underutilized, and to increase the production of ethanol from that particular saccharide source. In addition, based on increased ethanol production, the amount of saccharide source used in a fermentation reaction can be reduced, thereby saving costs associated with the saccharide source while producing the same amount of ethanol.

A number of complex saccharide substrates may be used as a starting source for depolymerization and subsequent fermentation using the cells and methods of the invention. Ideally, a recyclable resource may be used in the SSF process. Mixed waste office paper is a preferred substrate (Brooks *et al.*, (1995) *Biotechnol. Progress.* 11:619-625; Ingram *et al.*, (1995) U.S.P.N. 5,424,202), and is much more readily digested than acid pretreated bagasse (Doran *et al.*, (1994) *Biotech. Bioeng.* 44:240-247) or highly purified crystalline cellulose (Doran *et al.* (1993) *Biotechnol. Progress.* 9:533-538).

Glucanases, both endoglucanases and exoglucanases, contain a cellulose binding domain, and can be readily recycled for subsequent fermentations by harvesting the undigested cellulose residue using centrifugation (Brooks *et al.*, (1995) *Biotechnol. Progress.* 11:619-625). By adding this residue with bound enzyme as a starter, ethanol yields (per unit substrate) can be increased to over 80% of the theoretical yield with a concurrent 60% reduction in fungal enzyme usage. Such approaches work well with purified cellulose, although the number of recycling steps may be limited with substrates with a higher lignin content. Other substrate sources that are within the scope of the invention include any type of processed or unprocessed plant material, *e.g.*, lawn clippings, husks, cobs, stems, leaves, fibers, pulp, hemp, sawdust, newspapers, *etc.*

This invention is further illustrated by the following examples, which should not be construed as limiting.

EXEMPLIFICATION

Materials and Methods

In general, the practice of the present invention employs, unless otherwise indicated, conventional techniques of chemistry, molecular biology, recombinant DNA technology, PCR technology, immunology, microbiology, or cell culture, which are

within the skill of the art and are explained in the literature. See, e.g., Sambrook, Fritsch and Maniatis, *Molecular Cloning: Cold Spring Harbor Laboratory Press* (1989); *DNA Cloning*, Vols. 1 and 2, (D.N. Glover, Ed. 1985); *PCR Handbook Current Protocols in Nucleic Acid Chemistry*, Beaucage, Ed. John Wiley & Sons (1999); *Antibodies: A*

- 5 *Laboratory Manual*, Harlow et al., C.S.H.L. Press, Pub. (1999); *Bergey's Manual of Determinative Bacteriology*, Kreig et al., Williams and Wilkins (1984), and *Current Protocols in Molecular Biology*, eds. Ausubel et al., Wiley Interscience (1998).

For additional techniques for using host cells in various industrial applications including a fermentation reaction for producing, e.g., ethanol, see, e.g., Barrios-

- 10 Gonzalez et al., *Biotechnol. Ann. Rev.* 2:85-121 (1996); *From Ethnomycology to Fungal Biotechnology: Exploiting from Natural Resources for Novel Products*, Singh, J., Ed., Plenum Press, Pub. (1999); *Manual of Industrial Microbiology and Biotechnology*, Demain, A. Ed., Am. Soc. of Microbiology, Pub., (1999); *Biomining: Theory, Microbes, and Industrial Processes*, Rawlings, Ed., R.G. Landes Co., Pub. (1997);
- 15 *Biotechnology of Industrial Antibiotics*, Vandamme, E., Ed., Marcel Dekker, Pub. (1984); *Industrial Biotechnology*, Malik, V., Ed., Science, Pub. (1992); *Biotechnology and Food Ingredients*, Goldberg et al., Ed., Aspen Publishers (1991); *Biotechnology and Food Process Engineering*, Schwartzberg et al., Ed., Marcel Dekker, Pub. (1990); and *Food Biotechnology: Techniques and Applications*, Mittal, G., Technomic Pub. Co.
- 20 (1992).

Unless otherwise stated, the following materials and methods were used in the example that follows.

Fermentation media

- 25 Two types of media were tested: 1% CSL and Luria broth (LB). Each contained 97 g of either xylose or glucose per liter. Luria Broth consists of, per liter: 10 g Difco® Tryptone, 5 g Difco® Yeast Extract and 5 g NaCl. The 1% CSL medium consists of Corn Steep Liquor (1% w/v) plus mineral salts (mineral salts per liter: 1 g of KH_2PO_4 , 0.5 g of K_2HPO_4 , 3.1 g of $(\text{NH}_4)_2\text{SO}_4$, 0.4 g of $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$), and 20 mg $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$).

30

Media Preparation

Luria Broth was prepared by autoclaving a 2x nutrient stock containing, per liter: 20 g Difco® Tryptone, 10 g Difco® Yeast Extract, and 10 g NaCl. Sugars (xylose or glucose) were autoclaved separately. A 1% CSL medium was prepared as follows:

- 35 Commercial Corn Steep Liquor (CSL; 50% dry weight/50% water) was diluted with tap water to make a 20X stock containing 100 g dry weight of Corn Steep Liquor/L and was adjusted to pH 7.2 with NaOH (50%). After sterilization by autoclaving, the 20X CSL stock was clarified by centrifugation at 5000 x g for 10 minutes. Magnesium was added

as a 100X stock solution (40 g/L $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$). Iron was added as a 1000X stock solution made by dissolving 20 g $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ in 175 mL of HCl and adjusting to 1 L with sterile water. Nitrogen, sulfur and phosphorus were added using a 20X stock solution containing the following (per liter): 20 g of KH_2PO_4 , 10 g of K_2HPO_4 and 62 g of $(\text{NH}_4)_2\text{SO}_4$. Stock solutions containing minerals were sterilized separately by autoclaving. Final media was prepared by adding 750 mL of water to 1 L bottle containing 100 g of xylose or glucose. This solution was autoclaved, and stock solutions were added to provide a 1X concentration, and the final volume adjusted to 1 L. This broth was further diluted by the addition of supplements (acetaldehyde stock or distilled water) during fermentation with a resulting sugar content of 97 g/L.

Inoculum and Fermentation

Inocula were grown in same media used for fermentation and were prepared as follows. Three colonies from a fresh plate were used to inoculate seed cultures: 250-mL flask containing 100 mL medium. These cultures were grown with agitation (120 rpm) for 12-16 hours at 35°C to a final $\text{OD}_{550\text{nm}}$ of approximately 2.0-2.5. Cells were harvested by centrifugation at 5000 x g for 5 minutes, and resuspended in fresh media to an $\text{OD}_{550\text{nm}}$ of 0.1. The resulting suspension was distributed into 500-mL beakers containing 345 ml each of fermentation medium. Acetaldehyde was added as indicated. Batch fermentations (35 °C, 100 rpm) were maintained at pH 6.5 by the automatic addition of 2N KOH. Cell mass, ethanol, and base addition (2N KOH) were recorded during 96 hours of incubation.

Addition of the Nutrient Compound Acetaldehyde

A fresh stock solution was prepared containing 35 g/L acetaldehyde in water. Three methods of acetaldehyde supplementation were investigated: 1) single additions of acetaldehyde at the time of inoculation; 2) addition of acetaldehyde at the time of inoculation followed by a second addition after 12 hours of fermentation for 1% CSL with xylose or after 8 h for 1% CSL with glucose and LB with xylose; and 3) addition of acetaldehyde (1% CSL with xylose medium only) at the time of inoculation followed by subsequent additions after 12 h and 24 h. Equivalent amounts of distilled water were added instead of the acetaldehyde in control experiments.

Other Nutrient Additives

Additives (e.g., as listed in Table 1) were dissolved in 5mL di-H₂O. Solutions with pH lower than 5 were neutralized to pH 6.5 with 2N KOH. Nutritional supplements were then filter-sterilized (0.45 μ m filter disk) directly into the fermentation vessel containing the culture. The final concentration of all additives in the culture medium was 2 g/L unless otherwise indicated.

Analytical Procedures

Cell mass was estimated by measuring OD_{550nm} using Baush & Lomb® Spectronic 70 spectrophotometer. Based on experimental determinations, 1 ml of cell suspension at 1.0 OD was found to contain 0.33 mg of cell dry weight. Ethanol was measured by gas chromatography using a Varian® 3400 CX gas chromatograph with 1-propanol as an internal standard.

EXAMPLE 1

Methods for Improved Alcohol Production and Cell Growth in *Escherichia*

This example describes the effect of the addition of acetaldehyde to Luria broth (LB) and Xylose medium, 1% CSL and Xylose medium, and 1% CSL and glucose medium which have been inoculated with ethanologenic cells, i.e., recombinant *Escherichia coli* KO11. The effect of acetaldehyde on the production of ethanol and the growth of the ethanologenic cell, is described for each type of medium.

Tables 1, 3, and 5 show time to completion of fermentation, ethanol production, and volumetric productivity of the fermentation for all concentrations of the compound and the control. Maximum ethanol production is shown in grams/liter (g/L). The ethanol yield is shown in grams of ethanol/grams of added sugar. Maximum volumetric productivity is shown in grams ethanol/(liter)(hour). The average productivity is calculated from the start of fermentation to the completion of fermentation is grams ethanol/(liter)(hour).

Tables 2, 4, and 6 show the initial growth of the cells (OD taken after the first 24 hours of fermentation) and the maximum cell density (the time which culture reaches the maximum OD value).

Culture Results Using 1% CSL and Xylose

Figure 1, panel A shows ethanol production in g/L over a period of 96 hours.

Figure 1, panel B shows cell mass (OD 550nm), over 96 hours.

Figure 1 and Table 1 show the results of the addition of acetaldehyde to 1% CSL and Xylose medium over a time period of 96 hours compared to a control (no addition of acetaldehyde). Acetaldehyde was added to the fermentation medium at five different

- concentrations and at five different time points. Where only one acetaldehyde value is provided, this amount of acetaldehyde was added at the start of fermentation only (*i.e.*, 0.1 g/L acetaldehyde, 0.25 g/L acetaldehyde, and 0.5 g/L acetaldehyde). Where two additions are indicated, the first volume was added at the start of fermentation and the
- 5 second was added after 12 hours of fermentation. For example, 2x 0.25 g/L acetaldehyde refers to the addition of .25 g/L at the start of fermentation and .25 g/L after 12 hours. Where three additions are indicated, one was added at the start of fermentation, one after 12 hours and the third after 24 hours of fermentation. For example, 3x 0.25 g/L refers to the addition of 0.25 g/L at the start, 0.25 g/L after 12
- 10 hours and 0.25g/L after 24 hours.

Table 1. Culture Results Using 1% CSL Medium and Xylose (100g/L)

	Replicates	Time to completion	Ethanol Production		Ethanol yield	Volumetric Productivity		
			Maximum	%		Maximum	%	Average
Control	11	> 96 h	30.08	100.00	0.310	0.368	100.000	0.313
0.1g/L Acetaldehyde	2	> 96 h	29.51	98.11	0.304	0.389	103.594	0.307
0.25 g/L Acetaldehyde	2	> 96 h	34.45	114.53	0.335	0.425	115.326	0.359
0.5 g/L Acetaldehyde	2	> 96 h	42.66	141.84	0.439	0.623	169.163	0.444
2x 0.25 g/L Acetaldehyde	2	> 96 h	45.57	151.50	0.469	0.740	200.842	0.475
3x 0.25 g/L Acetaldehyde	2	> 96 h	38.21	127.03	0.393	0.540	146.543	0.398
Citrate	2	> 96 h	28.80	95.73	0.296	0.339	91.967	0.300
IsoCitrate	2	> 96 h	33.02	109.77	0.340	0.435	117.980	0.344
Glutamate	3	96	43.14	143.43	0.444	0.665	180.459	0.449
Glutamate + 0.25 Acetaldehyde	1	72	46.54	154.70	0.479	0.867	235.201	0.646
Glutamate 0.0.5 Acetaldehyde	1	72	44.97	149.51	0.463	0.818	221.931	0.625
Oxaloacetate	2	> 96 h	23.61	78.49	0.243	0.270	73.268	0.246
Aspartate	2	> 96 h	27.90	92.74	0.287	0.304	82.392	0.291
Pyruvate + 0.25 Acetaldehyde	2	72	45.71	151.94	0.470	0.920	249.715	0.635
Pyruvate + 0.5 Acetaldehyde	2	72	45.16	150.15	0.465	0.775	210.311	0.627
Pyruvate	5	72	44.42	147.66	0.457	0.792	215.013	0.617
Alanine	2	> 96 h	21.66	72.02	0.223	0.240	65.176	0.226
4g Pyruvate	1	72	44.80	148.94	0.461	0.838	227.413	0.622
1g Pyruvate	1	> 96 h	28.92	96.13	0.298	0.303	82.271	0.301
0.5g Pyruvate	1	> 96 h	29.24	97.22	0.301	0.324	87.855	0.305
0.25g Pyruvate	1	> 96 h	29.21	97.11	0.301	0.341	92.601	0.304
A-KG + Succinate	2	72	41.78	138.89	0.430	0.802	217.601	0.580
α-KG	2	72	42.32	140.70	0.436	0.866	234.952	0.588
Succinate	3	> 96 h	35.00	116.35	0.360	0.400	108.536	0.365
Fumarate + Malate	1	> 96 h	33.13	110.14	0.341	0.399	108.247	0.345
Yeast Extract	2	72	44.07	146.51	0.454	0.643	174.569	0.612
Casimino Acids	2	72	44.95	149.42	0.463	0.671	182.180	0.624
Fumarate	2	> 96	28.49	94.73	0.293	0.345	93.682	0.297
Malate	2	> 96	24.43	81.22	0.252	0.274	74.453	0.255

Table 2. Culture Results Using 1% CSL Medium and Xylose (100g/L)

	Replicates	Initial Growth (OD @ 24h)		Maximum Cell Density		
		OD	% Control	Time (h)	OD	% Control
Control	11	2.04	100.00	96	2.52	100.00
0.1g/L Acetaldehyde	2	2.25	110.04	48	2.56	101.92
0.25 g/L Acetaldehyde	2	2.48	121.20	96	3.12	124.15
0.5 g/L Acetaldehyde	2	2.55	124.64	96	4.31	171.13
2x 0.25 g/L Acetaldehyde	2	3.36	164.64	96	4.67	185.55
3x 0.25 g/L Acetaldehyde	2	3.04	148.81	48	4.02	159.74
Citrate	2	2.12	104.03	96	2.47	98.09
Isocitrate	2	2.17	106.01	96	3.01	119.43
Glutamate	3	3.69	180.49	96	5.04	200.16
Glutamate +0.25 Acetaldehyde	1	4.03	197.30	96	6.13	243.62
Glutamate 0.5 Acetaldehyde	1	4.10	200.52	96	5.98	237.83
Oxaloacetate	2	1.65	80.80	96	2.26	89.89
Aspartate	2	1.76	86.27	72	2.45	97.20
Pyruvate + 0.25 Acetaldehyde	2	4.28	209.46	72	6.19	245.84
Pyruvate + 0.5 Acetaldehyde	2	2.89	141.25	72	5.62	223.18
Pyruvate	5	4.11	201.20	96	5.98	237.58
Alanine	2	1.66	81.34	48	1.73	68.82
4g Pyruvate	1	5.00	244.69	72	6.58	261.52
1g Pyruvate	1	2.12	103.56	72	2.69	106.93
0.5g Pyruvate	1	2.05	100.14	72	2.64	104.79
0.25g Pyruvate	1	1.98	97.04	96	2.38	94.41
A-KG + Succinate	2	3.52	172.27	72	5.52	219.40
a-KG	2	3.55	173.77	72	5.66	224.85
Succinate	3	2.11	103.40	96	3.12	124.13
Fumarate + Malate	1	1.83	89.38	96	2.68	106.70
Yeast Extract	2	3.75	183.56	96	4.83	192.03
Casamino Acids	2	4.31	210.82	96	5.34	212.03
Fumarate	2	1.86	91.16	96	2.60	103.15
Malate	2	1.71	83.93	96	2.42	96.31

5

As shown in Tables 1 and 2, the addition of .25 g/L acetaldehyde at the start of fermentation and after 12 hours of fermentation in 1% CSL and xylose resulted in the highest amount of ethanol production as compared to the control. In addition, this concentration of acetaldehyde also resulted in the greatest initial growth and maximum cell density as compared to a control.

10

Culture Results Using 1% CSL and Glucose

Figure 2A shows ethanol production in g/L over a period of up to 96 hours.

Figure 2B shows cell mass (OD 550nm), over 96 hours.

- Figure 2 and Table 3 show the results of the addition of acetaldehyde to 1% CSL and glucose medium over a time period of up to 96 hours compared to a control (no addition of acetaldehyde). Acetaldehyde was added to the fermentation medium at five different concentrations and time points. Where only one acetaldehyde value is provided, this amount of acetaldehyde was added at the start of fermentation only (*i.e.*, 0.25 g/L acetaldehyde, 0.5 g/L acetaldehyde, and 0.75 g/L acetaldehyde). Where two additions are indicated, the first volume was added at the start of fermentation and the second was added after 8 hours of fermentation. For example, 0.5 + 0.25 g/L acetaldehyde refers to the addition of 0.5 g/L at the start of fermentation and 0.25 g/L after 8 hours, and 0.5 + 0.5/L acetaldehyde refers to the addition of 0.5 g/L at the start of fermentation and 0.5 g/L after 8 hours.

15

Table 3. Culture Results Using 1% CSL Medium and Glucose (100g/L)

	Replicates	Time to completion	Ethanol Production		Ethanol yield	Volumetric Productivity		Average	% Control
			Maximum	% Control		Maximum	% Control		
Control	13	96	36.99	100.00	0.381	0.514	100.00	0.385	100.00
0.25 g/L Acetaldehyde	1	72	40.31	108.97	0.415	0.678	132.01	0.560	145.29
0.5 g/L Acetaldehyde	3	96	39.95	107.99	0.411	0.615	119.73	0.416	107.99
0.75 g/L Acetaldehyde	1	96	43.44	117.43	0.447	0.699	136.11	0.452	117.43
0.5 + 0.25 g/L Acetaldehyde	1	96	40.25	108.82	0.414	0.598	116.48	0.419	108.82
0.5 + 0.5 g/L Acetaldehyde	1	96	40.03	108.22	0.412	0.659	128.25	0.417	108.22
Aspartate	1	72	41.79	112.96	0.430	0.759	147.72	0.580	150.61
Oxaloacetate (max vp @ 24)	1	72	39.72	107.37	0.409	0.712	138.48	0.552	143.16
Citrate	2	96	32.87	88.86	0.338	0.502	97.77	0.342	88.86
Isocitrate	2	72	33.95	91.77	0.349	0.588	114.38	0.471	122.36
Glutamate (max vp @ 24)	1	72	43.39	117.30	0.447	0.763	148.52	0.603	156.40
a-ketoglutarate	2	72	40.33	109.03	0.415	0.657	127.95	0.560	145.37
Pyruvate (max vp @ 24)	4	96	42.35	114.48	0.436	0.663	129.06	0.441	114.48
Alanine (max vp @ 24)	2	96	34.37	92.90	0.354	0.468	91.10	0.338	92.90
4g Pyruvate (max vp @ 24)	2	96	44.83	121.18	0.461	0.585	113.84	0.467	121.18
Succinate (max vp @ 24)	2	96	30.13	81.46	0.310	0.511	99.45	0.314	81.46
Fumarate (max vp @ 24)	2	96	32.92	88.98	0.339	0.559	108.88	0.343	88.98
Malate (max vp @ 24)	2	96	27.58	74.56	0.284	0.498	96.97	0.287	74.56

Stimulation with glucose was smaller than xylose, but present. Oxaloacetate stimulated more with glucose, while 2-ketoglutarate stimulated more in xylose

Table 4. Culture Results Using 1% CSL Medium and Glucose (100g/L)

	Replicates	Initial Growth (OD@24h)		Maximum Cell Density		
		OD	% Control	Time (h)	OD	% Control
Control	13	2.67	100.00	48	2.69	100.00
0.25 g/L Acetaldehyde	1	3.00	112.34	72	3.17	118.01
0.5 g/L Acetaldehyde	3	2.77	103.55	72	2.96	110.00
0.75 g/L Acetaldehyde	1	2.47	92.47	72	2.81	104.56
0.5 + 0.25 g/L Acetaldehyde	1	2.64	98.90	48	2.99	111.40
0.5 + 0.5 g/L Acetaldehyde	1	2.74	102.52	48	3.09	115.00
Aspartate	1	2.86	107.10	72	3.79	140.94
Oxaloacetate (max vp @ 24)	1	3.29	123.05	72	3.46	128.90
Citrate	2	2.43	90.97	72	2.58	95.99
Isocitrate	2	2.78	103.82	72	2.62	97.64
Glutamate (max vp @ 24)	1	2.86	107.10	72	3.79	140.94
α -ketoglutarate	2	3.29	123.05	72	3.46	128.90
Pyruvate (max vp @ 24)	4	3.46	129.39	72	3.83	142.47
Alanine (max vp @ 24)	2	2.08	77.97	96	2.68	99.80
4g Pyruvate (max vp @ 24)	2	3.84	143.44	72	4.07	151.63
Succinate (max vp @ 24)	2	2.69	100.47	72	2.66	98.99
Fumarate (max vp @ 24)	2	2.67	99.76	72	3.17	117.92
Malate (max vp @ 24)	2	3.19	119.49	72	3.39	126.18

- As shown in Table 4, addition of .75 g/L of acetaldehyde at the start of fermentation in 1% CSL and glucose resulted in the highest amount of ethanol production as compared to the control. The addition of 0.25 g/L of acetaldehyde at the start of fermentation also resulted in the greatest initial growth and maximum cell density as compared to a control.

10 Culture Results Using LB and Xylose

Figure 3A shows ethanol production in g/L over 72 hours. Figure 3B shows cell mass (OD 550nm), over 72 hours.

- Figure 3 and Table 5 show the results of the addition of acetaldehyde to LB and xylose medium over a time period of 72 hours as compared to a control (no addition of acetaldehyde). Acetaldehyde was added to the fermentation medium at five different concentrations and time points. Where only one acetaldehyde value is provided, this amount of acetaldehyde was added at the start of fermentation only (*i.e.*, 0.25 g/L acetaldehyde, 0.5 g/L acetaldehyde, and 0.75 g/L acetaldehyde). Where two additions are indicated, the first volume was added at the start of fermentation and the second was added after 8 hours of fermentation. For example, 0.5 + 0.25 g/L acetaldehyde refers to the addition of 0.5 g/L at the start of fermentation and 0.25 g/L after 8 hours, and 0.5 + 0.5/L acetaldehyde refers to the addition of 0.5 g/L at the start of fermentation and 0.5 g/L after 8 hours.

Table 5. Culture Results Using Luria Broth Medium and Xylose (100g/L)

	Replicates	Time to completion	Ethanol Production		Ethanol yield	Volumetric Productivity		Average	% Control
			Maximum	%		Maximum	%		
Control	5	48	43.08	100.00	0.443	1.152	100.00	0.897	100.00
0.25 g/L Acetaldehyde	1	48	43.80	101.68	0.451	1.352	117.36	0.913	101.68
0.5 g/L Acetaldehyde	2	48	42.59	98.88	0.438	1.027	89.20	0.887	98.88
0.75 g/L Acetaldehyde	1	48	42.65	99.00	0.439	0.916	79.50	0.889	99.00
0.5 + 0.25 g/L Acetaldehyde	2	48	41.40	96.11	0.426	0.904	78.52	0.863	96.11
0.5 + 0.5 g/L Acetaldehyde	2	48	41.64	96.65	0.429	0.890	77.26	0.867	96.65
α -ketoglutarate	2	48	42.47	98.60	0.437	1.348	117.02	0.885	98.60
Glutamate	2	48	44.67	103.69	0.460	1.328	115.30	0.931	103.69
2g Pyruvate	2	48	44.35	102.96	0.457	1.284	111.47	0.924	102.96
4g Pyruvate	2	48	45.13	104.76	0.465	1.166	101.27	0.940	104.76

Table 6. Culture Results Using Luria Broth Medium and Xylose (100g/L)

	Replicates	Initial Growth (OD at 24 hours)		Maximum Cell Density		
		OD	% Control	Time (h)	OD	% Control
Control	5	9.99	100.00	48	11.04	100.00
0.25 g/L Acetaldehyde	1	11.07	110.79	24	11.07	100.20
0.5 g/L Acetaldehyde	2	10.86	108.77	48	11.92	107.97
0.75 g/L Acetaldehyde	1	10.54	105.54	48	12.25	110.94
0.5 + 0.25 g/L Acetaldehyde	2	9.89	99.02	48	10.81	97.92
0.5 + 0.5 g/L Acetaldehyde	2	10.68	106.98	48	11.78	106.68
a-ketoglutarate	2	10.27	102.79	48	10.59	95.90
Glutamate	2	10.61	106.21	24	10.61	96.07
2g Pyruvate	2	11.30	113.09	24	11.30	102.29
4g Pyruvate	2	11.45	114.64	48	11.65	105.48

As shown in Table 3, addition of 0.5 g/L of acetaldehyde at the start of fermentation in LB + xylose resulted in the highest amount of ethanol production as compared to the control. The addition of 0.5 g/L of acetaldehyde at the start of fermentation results in the greatest initial growth and maximum cell density as compared to the control.

EXAMPLE 2

Methods for Improved Alcohol Production and Cell Growth in *Klebsiella*

This example describes the effect of the addition of acetaldehyde to 1% CSL and xylose medium, which has been inoculated with ethanologenic, cells, *i.e.*, *Klebsiella oxytoca* P2. The effect of acetaldehyde on the production of ethanol and the growth of the ethanologenic cell is presented in Table 7.

More particularly, Table 7 shows time to completion of fermentation, ethanol production and volumetric productivity of the fermentation for all concentrations of the compound and the control. Maximum ethanol production is shown in grams/liter (g/L). The ethanol yield is shown in grams of ethanol/grams of added sugar. Maximum volumetric productivity is shown in grams of ethanol/(liter)(hour). The average productivity is calculated from the start of fermentation to the completion of fermentation in grams ethanol/(liter)(hour). Table 7 also shows the initial growth of the cells (OD taken after the first 24 hours of fermentation) and the maximum cell density (the time which the culture reaches the maximum OD value at 48 hours).

Importantly, Table 7 shows the results of the addition of acetaldehyde to 1% CSL and xylose medium over a time period of 72 to 96 hours compared to a control (no addition of acetaldehyde) using an inoculum of the ethanologenic host *Klebsiella oxytoca* P2. Acetaldehyde was added to the fermentation medium at five different

concentrations and time points. Where only one acetaldehyde value is provided, this amount of acetaldehyde was added at the start of fermentation only (*i.e.*, 0.25 g/L acetaldehyde or 0.5 g/L). Where two or three additions are indicated, the first volume was added at the start of fermentation and additional doses of the nutrient compound

5 were added after 8 hours of fermentation or at 8-hour intervals.

As shown in Table 7, addition of 0.5 g/L of acetaldehyde at the start of fermentation in CSL and xylose resulted in the highest amount of ethanol production as compared to the control. The addition of 0.5 g/L of acetaldehyde at the start of fermentation also resulted in increased initial growth and maximum cell density as

10 compared to the control.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
21

Table 7. Culture Results Using 1% CSL Medium and Xylose (100g/L)

	Time to completion	Ethanol Production				Volumetric Productivity			
		Maximum	% Control	Ethanol yield	Maximum	% Control (based on Max)	Average	% Control (based on Avg)	
Control	> 96 h	23.69	100.00	0.244	0.34	100.00	0.247	100.00	
0.25 g/L Acetaldehyde	> 96 h	25.43	107.35	0.262	0.47	135.46	0.265	107.35	
0.5 g/L Acetaldehyde	> 96 h	26.53	111.98	0.273	0.51	147.11	0.276	111.98	
2x 0.25 g/L Acetaldehyde	> 96 h	25.56	107.88	0.263	0.44	126.68	0.266	107.88	
3x 0.25 g/L Acetaldehyde	> 96 h	26.14	110.32	0.269	0.47	135.55	0.272	110.32	

	Initial Growth (OD @ 24h)		Maximum Cell Density	
	OD	% Control	Time (h)	% Control
Control	3.77	100.00	72.00	100.00
0.25 g/L Acetaldehyde	3.54	93.90	72.00	5.68
0.5 g/L Acetaldehyde	2.61	69.39	96.00	5.89
2x 0.25 g/L Acetaldehyde	3.34	88.59	96.00	5.30
3x 0.25 g/L Acetaldehyde	3.60	95.66	72.00	5.40

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following

5 claims. Moreover, any number of genetic constructs, host cells, and methods described in United States Patent Nos. 5,821,093; 5,482,846; 5,424,202; 5,028,539; 5,000,000; 5,487,989, 5,554,520, and 5,162,516, may be employed in carrying out the present invention and are hereby incorporated by reference.